

## ORIGINAL SCIENTIFIC PAPER

# Relationship between Tibia Length Measurements and Body Height: A Prospective Regional Study among Adolescents in the Eastern Region of Kosovo

Fitim Arifi<sup>1</sup>, Bojan Masanovic<sup>2</sup>, Jovan Gardasevic<sup>2</sup> and Dusko Bjelica<sup>2</sup>

<sup>1</sup>University of Tetovo, Faculty of Physical Education, Tetovo, North Macedonia, <sup>2</sup>University of Montenegro, Faculty for Sport and Physical Education, Niksic, Montenegro

## Abstract

This study aims to determine standing height in males and females from the Eastern Region in Kosovo and its association with tibia length, as an alternative to standing height prediction. A total of 364 individuals (185 male and 179 female) took part in this study. The anthropometric measurements were provided in accordance with the ISAK guidelines. The relationships between body height and tibia length were defined using simple correlation coefficients at a ninety-five per cent confidence interval. A means comparison in standing height and tibia length between males and females was performed using a t-test. In the end, a linear regression analysis was implemented to examine how tibia length can reliably predict standing height. The results showed that Eastern Kosovan males are  $178.79 \pm 6.07$  cm tall and have a tibia length of  $38.58 \pm 2.49$  cm, while Eastern Kosovan females are  $164.74 \pm 4.89$  cm tall and have a tibia length of  $35.32 \pm 1.85$  cm. This finding defines Eastern Kosovans, male and female, as a tall group, but slightly shorter than the general Kosovan population. Moreover, tibia length reliably predicts standing height in Kosovan males and females but not as sufficiently reliably as arm span does. This study also indicates the need to develop separate height models for each region in Kosovo because the results from Eastern Kosovans do not correspond to general values.

**Keywords:** prediction, measurement, stature, tibia length, Kosovan

## Introduction

Kosovo is located in south-eastern Europe, in the central region of the Balkan Peninsula. Covering an area of 10,908 km<sup>2</sup>, it has a population of 1,883,018 inhabitants (Gardasevic, 2019a; Arifi, Gardasevic, & Masanovic, 2018). Administratively, it is subdivided into five regions (Eastern, Western, Northern, Southern, and Central) and is bordered by four countries (Albania, Macedonia, Montenegro, and Serbia) (Arifi et al., 2017). This study aims to analyse the standing height of adolescents from the Eastern region of Kosovo and its estimation utilizing tibia length measurements. This region contains two districts (Ferizaj and Gjiilan) and eleven municipalities (Ferizaj, Hani i Elezit, Kaçanik, Štimlje/Shtime, Štrpce/Shtërpçë, Gjiilan,

Kamenica, Klokot, Partesh, Ranilug, and Vitina). Covering an area of 2,236 km<sup>2</sup>, it has a population of 366,589 inhabitants (Masanovic, Gardasevic, & Arifi, 2018; Gardasevic, 2019a; Arifi, Masanovic, & Gardasevic, 2020). While Kosovo is small, it has a highly varied terrain. One part belongs in the Dinaric Alps, so it can be assumed this fact might influence this study's main objective because of the type of the soil and other socio-economic and geographic characteristics as potential influencing factors (Popovic, 2018).

Many scientific studies confirm that the measurement of standing height is an essential element in assessing nutritional status and evaluating basic energy requirements (Arifi et al., 2017; Datta Banik, 2011; Popovic, Arifi, & Bjelica, 2017). It is



Correspondence:

B. Masanovic  
University of Montenegro, Faculty for Sport and Physical Education, Narodne omladine bb, 81400, Niksic, Montenegro  
E-mail: bojanma@ucg.ac.me

also a significant factor in assessing children's growth, adjusting the measures of physical capacity, predicting drug dosages and setting standards of physiological variables, such as muscle strength, metabolic rate, lung volume, and glomerular filtration (M. Golshan, Crapo, Amra, Jensen, & R. Golshan, 2007; Ter Goon, Toriola, Musa, & Akusu, 2011; Gardasevic, Rasidagic, Krivokapic, Corluka, & Bjelica, 2017).

However, exact standing height is sometimes impossible to measure in the standard way (e.g., due to paralysis, fractures, amputation, scoliosis, and pain). In such situations, standing height estimation should be assessed from other reliable anthropometric indicators, such as hand and tibia length, foot length, knee height, length of the forearm, length of the sternum, vertebral column length, sitting height, length of scapula, arm span as well as cranial sutures, skull, facial measurements et cetera (Fredriks et al., 2005; Popovic, Gardasevic, Masanovic, Arifi, & Bjelica, 2017). All these anthropometric indicators, which are used as an alternative to standing height prediction, are also crucial in assessing decreases in body height that emerge as a result of ageing. They are also very helpful when it is necessary to determine the standing height of a person with disproportionate growth abnormalities, skeletal dysplasia, mobility problems, kyphosis, in a person with standing height loss during surgical procedures on the spine as well to anticipate standing height in older people who cannot keep upright or maintain erect posture due to ageing (Hickson & Frost, 2003; Masanovic, Arifi, & Gardasevic, 2019). Lastly, it should not be forgotten that this knowledge is relevant in sport science because standing height represents a significant factor influencing success in various sport disciplines (Gusic, Popovic, Molnar, Masanovic, & Radakovic, 2017; Popovic, 2017; Arifi, Bjelica, & Masanovic, 2019; Masanovic, 2019).

Numerous researchers emphasize the benefit of using various body parameters in predicting standing height, and almost all agree that arm span is the most reliable one in adults (Mohanty, Babu, & Nair, 2001; Brown, Feng, & Knapp, 2002; Bjelica et al., 2012; Masanovic, 2018). Some authors believe that foot length measurement is the most reliable predictor during adolescence (Jakhar, Pal, & Paliwal, 2010; Popovic et al., 2017), because ossification occurs earlier in the foot than in the long bones, and standing height will be more accurately predicted from this measurement than by using long bones during adolescence.

Furthermore, it has been established that the relationship of long bones and standing height varies in different ethnic and racial groups (Steele & Chenier, 1990; Reeves, Varakamin, & Henry, 1996; Quanjer et al., 2014; Khatun, Sharma, Jain, & Gupta, 2016) and also in various regions (Milasinovic, Popovic, Matic, Gardasevic, & Bjelica, 2016; Masanovic, Gardasevic, & Arifi, 2018a; Masanovic, Bavcevic, & Prskalo, 2019; Masanovic, Arifi, & Gardasevic, 2020; Masanovic, Arifi, & Gardasevic, 2020a); therefore, researchers propose designing specific formulas for calculating standing height from long bones for each ethnic or racial group, and for different regions. The mentioned variations might also be the case with tibia length predictions, mostly be-

cause the Dinaric Alps population has a specific body composition (Popovic, 2017; Starc et al., 2020; Popovic, Masanovic, Martinovic, Bjelica, & Gardasevic, 2020). Many studies on this subject are available worldwide; some data is available on Kosovan subjects, but most have covered the whole Kosovan population, and few have studied regional differences. Some studies confirm that Western Kosovans, Southern Kosovans, and Northern Kosovans have specific standing height/tibia length ratios compared to the general population in Kosovo (Gardasevic, 2019b; Gardasevic, Masanovic, & Arifi, 2018; Gardasevic, Masanovic, & Arifi, 2019). Considering the rather sparse recent scientific literature, the purpose of this research was to examine the standing height in both Eastern Kosovan genders and its association with tibia length.

## Method

This research gathered 364 final-year high-school students (185 male and 179 female) from Kosovo's Eastern Region. The reasons that qualified the selected individuals are 1) related to the fact that the growth of an individual ceases by this age, and 2) related to the fact that there is no age-related loss in standing height at this age. The average age of the male subject was  $18.20 \pm 0.40$  years old (range 18–19 years), while the average age of the female subject was  $18.15 \pm 0.36$  years old (range 18–19 years). It is important to underline that the researchers have excluded from the data analysis the individuals with physical deformities and those without informed consent. Another exclusion criterion was being non-Eastern Kosovan.

The anthropometric measurements, including standing height and tibia length, were conducted according to the International Society for the Advancement of Kinanthropometry (ISAK) protocol (Marfell-Jones, Olds, Stew, & Carter, 2006). Trained measurers conducted the assessment of selected anthropometric indicators, while the quality of their performance was evaluated against the prescribed "ISAK Manual". Finally, the age of each subject was obtained directly from their birthdays.

The analysis was performed by using the Statistical Package for Social Sciences (SPSS) version 20.0 (Chicago, IL, USA) adjusted for use on personal computers. Means and standard deviations (SD) were obtained for both anthropometric variables. A comparison of means of standing height and tibia length between genders was performed using a t-test. The relationships between standing height and tibia length were determined using simple correlation coefficients at a ninety-five per cent confidence interval. Next, a linear regression analysis was carried out to examine the extent to which the tibia length can reliably predict standing height. Statistical significance was set at  $p < 0.05$ .

## Results

In Table 1, a summary of the anthropometric measurements in both genders is shown. The mean standing height for males was  $178.79 \pm 6.07$  centimetres, and tibia length was  $38.58 \pm 2.49$  centimetres, while for females the standing height was  $164.74 \pm 4.89$

**Table 1.** Anthropometric Measurements of the Study Subjects

Subjects	Body Height Range (Mean±Stand.Dev.)	Tibia Length Range (Mean± Stand.Dev.)
Male	161.4-193.0 (178.79±6.07)	30.1-45.7 (38.58±2.49)
Female	153.3-178.0 (164.74±4.89)	31.0-40.3 (35.32±1.85)

centimetres, and tibia length was 35.32±1.85 centimetres. The gender difference between standing height and tibia length measurements was significant (standing height: t=24.849; p<.000; and tibia length: t=14.177; p<.000).

In Table 2, the simple correlation coefficients and their

ninety-five per cent confidence interval analysis between the anthropometric measurements are displayed. The associations between standing height and tibia length were significant (p<0.000) and high in this sample, regardless of gender (male: 0.671; female: 0.640).

**Table 2.** Correlation between Body Height and Tibia Length of the Study Subjects

Subjects	Correlation Coefficient	95% confidence interval	Significance p-value
Male	0.671	0.562-0.779	<0.000
Female	0.640	0.526-0.754	<0.000


In Table 3, the results of the linear regression analysis are shown. The first of all models were extracted by including age as a covariate. However, it was found that the contribution of age was insignificant; therefore, it was eliminated as a factor from the study, and estimations were derived as a univariate analy-

sis. The high values of the regression coefficient (male: 0.671; female: 0.640) signify that tibia length notably predicts standing height in both Eastern Kosovan genders (male: t=12.227, p<0.000; female: t=11.072, p<0.000), which confirms the R-square (%) for the male (45.0) as well as for the female (40.9).

**Table 3.** Results of Linear Regression Analysis Where the Tibia Length Predicts the Body Height

Subjects	Regression Coefficient	Standard Error (SE)	R-square (%)	t-value	p-value
Male	0.671	4.514	45.0	12.227	0.000
Female	0.640	3.638	40.9	11.072	0.000

The associations between tibia length measurements and standing height among the above models are plotted as a scatter diagram (Figure 1).



**FIGURE 1.** Scatter Diagram and Relationship between Tibia Length Measurements and Body Height among Both Genders

**Discussion**

Over the centuries, many researchers have studied the assessment of standing height using various anthropometric measures. It must be emphasized that arm span has been evaluated as the most reliable body indicator for predicting standing height (Ter Goon et al., 2011; Popovic, Bjelica, Georgiev, Krivokapic, & Milasinovic, 2016; Masanovic, Popovic, Jarani, Spahi, & Bjelica, 2020), while tibia length was a very close second (Agnihotri, Smita, Jowaheer, & Pratap, 2009; A. Kaore, B.P. Kaore, Kamdi, & S. Kaore, 2012; Khatun et al., 2016). Furthermore, it should be considered that the individual and

ethnic variations referring to standing height and its association with tibia length can vary from among ethnic groups and races. Consequently, the effect of racial and ethnic differences on these measures reduces the possibility of generalization (Popovic, 2018; Masanovic et al., 2019). This study’s findings are consistent with results of the study conducted by Agnihotri et al. (2009), which confirmed a very high linear correlation between standing height and tibia length in both genders of an Indo-Mauritian population, and with the results of the study conducted by Khatun et al. (2016), which showed a significant correlation between standing height and tibia length in

both genders of an Indian population, in which the correlation coefficient for tibia length is  $r=0.67$  in males and  $r=0.58$  in females. As the correlation between tibia length and standing height was significant in both Eastern Kosovan genders, the tibia length measure, therefore, seems to be a reliable indirect anthropometric indicator for estimating standing height in both genders of the Eastern Kosovan population.

The necessity of developing separate standing height models for each population on account of ethnic differences has been confirmed by several studies. Arifi et al. (2017) analysed the entire Kosovan population and found specific correlation coefficients between standing height and arm span in Kosovan male ( $r=0.794$ ) and female ( $r=0.766$ ) population. Also, some recent studies have confirmed the regional differences between the same ethnic groups (Arifi, 2017; Popovic et al., 2017). Care should be taken here because the three regional studies indicate the specificity for height prediction using the tibia. In the first study, Gardasevic (2019b) found specific correlation coefficients for standing height and tibia length in Western Kosovan males ( $r=0.538$ ) and females ( $r=0.559$ ) compared with the entire Kosovan population; in the second study, Gardasevic et al. (2018) found specific correlation coefficients for standing height and tibia length in Southern Kosovan males ( $r=0.734$ ) and females ( $r=0.639$ ) compared with the entire Kosovan population; in the third study, Gardasevic et al. (2019) found specific correlation coefficients in standing height and tibia length in Northern Kosovan males ( $r=0.730$ ) and females ( $r=0.675$ ) compared with the entire Kosovan population. Therefore, the main goal of this study is to examine whether this hypothesis also applies to the Eastern Kosovans (i.e., for one the of five Kosovan regions). The results of the studies mentioned previously confirm the necessity

of developing separate standing height models for both genders in Kosovo, and also recommended that further studies should consider dividing the population of this country into regional subsamples and analysing them separately to remove the doubt that geographical differences exist (such as type of soil) influencing the average standing height in both Kosovan genders as well as its association with tibia length. This was based on the fact that the whole of Kosovo does not fall into the Dinaric Alps racial classification (Masanovic et al., 2019b). In parallel, this study confirms that assumption and also that it is necessary to develop separate standing height models for each population on account of regional variations in Kosovo.

Finally, it is necessary to point out that a limitation of this study might be the composition of the respondent sample, which comprised high school students. This limitation is based on the fact that some studies question whether the growth of an individual ceases by this age (Grasgruber et al., 2017). This assumption might be supported by the fact that university-educated individuals are taller than the high school graduate population in Hungary and Poland (Szollosi, 1998; Wronka & Pawlinska-Chmara, 2009). However, this was not the case in Montenegro (Popovic et al., 2017). This dilemma could be resolved by comparing the average standing height measures of this study to the results of a study with a sample of university students. Another limitation of this study is the fact that both genders of the subjects in Kosovo had not yet reached their full genetic potential, since various environmental factors controlled their development (war and the difficult economic situation of the previous three decades). Further continuous monitoring is necessary, mostly because the secular changes influencing standing height are expected to rise in the next two to three decades.

#### Acknowledgements

There are no acknowledgements.

#### Conflict of Interest

The authors declare that there are no conflicts of interest.

**Received:** 18 November 2020 | **Accepted:** 05 January 2021 | **Published:** 01 June 2021

#### References

- Agnihotri, A. K., Smita, K., Jowaheer, V., & Pratap, A. (2009). Estimation of stature from percutaneous length of tibia and ulna in Indo-Mauritian population. *Forensic Science International*, 187(1-3), 109.e1-109.e3, 2009. doi: 10.1016/j.forsciint.2009.02.010
- Arifi, F., Masanovic, B., & Gardasevic, J. (2020). Relationship between Sitting Height Measurements and Standing Height: A Prospective Regional Study among Adolescents in the Northern Region of Kosovo. *Sport Mont*, 18(3), 35-39. doi: 10.26773/smj.201012
- Arifi, F., Bjelica, D., & Masanovic, B. (2019). Differences in anthropometric characteristics among junior soccer and handball players. *Sport Mont*, 17(1), 45-49. doi: 10.26773/smj.190208
- Arifi, F., Gardasevic, J., & Masanovic, B. (2018). Relationship between foot length measurements and body height: A prospective regional study among adolescents in central region of Kosovo. *Sport Mont*, 16(3), 75-79.
- Arifi, F., Bjelica, D., Sermahaj, S., Gardasevic, J., Kezunovic, M., & Popovic, S. (2017). Stature and its estimation utilizing arm span measurements in Kosovan adults: national survey. *International Journal of Morphology*, 35(3), 1161-1167.
- Arifi, F. (2017). Stature and its estimation utilizing arm span measurements of both gender adolescents from southern region in Kosovo. *Sport Science*, 10(1), 92-95.
- Bjelica, D., Popovic, S., Kezunovic, M., Petkovic, J., Jurak, G., & Grasgruber, P. (2012). Body Height and Its Estimation Utilizing Arm Span Measurements in Montenegrin Adults. *Anthropological Notebooks*, 18(2), 69-83.
- Brown, J. K., Feng, J. Y., & Knapp, T. R. (2002). Is self-reported height or arm span a more accurate alternative measure of height? *Clinical Nursing Research*, 11(4), 417-432.
- Datta Banik, S. (2011). Arm span as a proxy measure for height and estimation of nutritional status: a study among Dhimals of Darjeeling in West Bengal India. *Annals of Human Biology*, 38(6), 728-735.
- Fredriks, A. M., Van Buuren, S., van Heel, W. J. M., Dijkman-Neerick, R. H. M., Verloove-Vanhorich, S. P., & Wit, J. M. (2005). Nationwide age references for sitting height, leg length, and sitting height/height ratio, and their diagnostic value for disproportionate growth disorders. *Archives of Disease in Childhood*, 90(8), 807-812, 2005.
- Gardasevic, J. (2019a). Body height in Kosovo population and its estimation from tibia length: National survey. *Anthropological Notebooks*, 25(3), 77-86.
- Gardasevic, J. (2019b). Standing Height and its Estimation Utilizing Tibia Length Measurements in Adolescents from Western Region in Kosovo. *International Journal of Morphology*, 37(1), 227-231.
- Gardasevic, J., Masanovic, B., & Arifi, F. (2019). Relationship between Tibia Length Measurements and Standing Height: A Prospective Regional Study among Adolescents in Northern Region of Kosovo. *Anthropologie*, 57(3), 263-269. doi: 10.26720/anthro.19.08.28.2
- Gardasevic, J., Masanovic, B., & Arifi, F. (2018). Relationship between tibia length measurements and standing height: A prospective regional study among adolescents in southern region of Kosovo. *Sport Mont*, 16(3), 51-55. doi: 10.26773/smj.181009
- Gardasevic, J., Rasidagic, F., Krivokapic, D., Corluca, M., & Bjelica, D. (2017). Stature and Its Estimation Utilizing Arm Span Measurements in Male Adolescents from Federation of Bosnia and Herzegovina Entity in Bosnia and Herzegovina. *Montenegrin Journal of Sports Science and Medicine*, 6(1), 37-44.
- Golshan, M., Crapo, R. O., Amra, B., Jensen, R. I., & Golshan, R. (2007). Arm span as an independent predictor of pulmonary function parameters: validation and reference values. *Respirology*, 12(3), 361-366.
- Grasgruber, P., Popovic, S., Bokuvka, D., Davidovic, I., Hřebickova, S., Ingrova, P., Potpara, P., Prce, S., & Stracarova, N. (2017). The mountains of giants: an anthropometric survey of male youths in Bosnia and Herzegovina. *Royal Society Open Science*, 4, 161054.
- Gusic, M., Popovic, S., Molnar, S., Masanovic, B., & Radakovic, M. (2017).

- Sport-Specific Morphology Profile: Differences in Anthropometric Characteristics among Elite Soccer and Handball Players. *Sport Mont*, 15(1), 3-6.
- Hickson, M., & Frost, G. A. (2003). Comparison of three methods for estimating height in the acutely ill elderly population. *Journal of Human Nutrition and Dietetics*, 16(1), 13-20.
- Jakhar, J. K., Pal, V., & Paliwal, P. K. (2010). Estimation of height from measurements of foot length in Haryana region. *Journal of Indian Academy of Forensic Medicine*, 32(3), 231-233.
- Kaore, A., Kaore, B. P., Kamdi, A., & Kaore, S. (2012). Stature estimation from tibial length. *National Journal of Integrated Research in Medicine*, 3(2), 51-56.
- Khatun, S. S., Sharma, N., Jain, S. K., & Gupta, A. (2016). Estimation of stature from percutaneous tibial length in Indian population. *International Journal of Anatomy and Research*, 4(3), 2571-2576. doi: 10.16965/ijar.2016.278
- Marfell-Jones, M., Olds, T., Stew, A. D., & Carter, J. E. L. (2006). *International standards for anthropometric assessment*. Potchesfroom: International Society for the Advancement of Kinanthropometry.
- Masanovic, B., Popovic, S., Jarani, J., Spahi, A., & Bjelica, D. (2020). Nationwide stature estimation from armspan measurements in Albanian youngsters. *International Journal of Morphology*, 38(2), 382-388.
- Masanovic, B., Arifi, F., & Gardasevic, J. (2020). Standing Height Estimation from Sitting Height Measurements in Adolescents in the Central Region of Kosovo. *Sport Mont*, 18(2), 19-23.
- Masanovic, B., Arifi, F., & Gardasevic, J. (2020a). Relationship between sitting height measurements and standing height: A prospective regional study among adolescents in the southern region of Kosovo. *International Journal of Morphology*, 38(6), 1681-1685.
- Masanovic, B. (2019). Comparative Study of Morphological Characteristics and Body Composition between Different Team Players from Serbian Junior National League: Soccer, Handball, Basketball and Volleyball. *International Journal of Morphology*, 37(2), 612-619.
- Masanovic, B., Gardasevic, J., & Arifi, F. (2019). Relationship between foot length measurements and body height: a prospective regional study among adolescents in northern region of Kosovo. *Anthropologie*, 57(2), 227-233, 2019. doi: 10.26720/anthro.18.01.23.1.
- Masanovic, B., Arifi, F., & Gardasevic, J. (2019). Standing height and its estimation utilizing sitting height measurements in adolescents from the western region in Kosovo. *Sport Mont*, 17(3), 3-7.
- Masanovic, B., Bavcevic, T., & Prskalo, I. (2019). Regional differences in adult body height in Kosovo. *Montenegrin Journal of Sports Science and Medicine*, 8(1), 69-76.
- Masanovic, B. (2018). Standing height and its estimation utilizing arm span and foot length measurements in dinaric alps population: a systematic review. *Sport Mont*, 16(2), 101-106.
- Masanovic, B., Gardasevic, J., & Arifi, F. (2018). Relationship between Foot Length Measurements and Body Height: A Prospective Regional Study among Adolescents in Eastern Region of Kosovo. *Sport Mont*, 16(1), 9-13. doi: 10.26773/smj.180202
- Masanovic, B., Gardasevic, J., & Arifi, F. (2018a). Relationship between foot length measurements and standing height: a prospective regional study among adolescents in southern region of Kosovo. *Sport Mont*, 16(2), 27-31. doi: 10.26773/smj.180605
- Milasinovic, R., Popovic, S., Matic, R., Gardasevic, J., & Bjelica, D. (2016). Body Height and its Estimation Utilizing Arm Span Measurements in Male Adolescents from Southern Region in Montenegro. *Sport Mont*, 14(2), 21-23.
- Mohanty, S. P., Babu, S. S., & Nair, N. S. (2001). The use of arm span as a predictor of height. A study of South Indian women. *Journal of Orthopaedic Surgery (Hong Kong)*, 9(1), 19-23.
- Popovic, S., Masanovic, B., Martinovic, S., Bjelica, D., & Gardasevic, J. (2020). Trajectories in Body Height, Body Weight, BMI, and Nutrition Status from 1979 to 1987: A Measurement-Based Analysis of 15,717 Male Adolescents from the Capital City of Montenegro. *Frontiers in Public Health* 8, 610358. doi: 10.3389/fpubh.2020.610358
- Popovic, S. (2018). Arm-span measurement as an alternative estimation of true height in Montenegrin young adults of both sexes: A national survey. *Anthropological Notebooks*, 24(1), 53-67.
- Popovic, S. (2017). Local Geographical Differences in Adult Body Height in Montenegro. *Montenegrin Journal of Sports Science and Medicine*, 6(1), 81-87.
- Popovic, S., Arifi, F., & Bjelica, D. (2017). Standing Height and its Estimation Utilizing Foot Length Measurements in Kosovan Adults: National Survey. *International Journal of Applied Exercise Physiology*, 6(2), 1-7.
- Popovic, S., Gardasevic, J., Masanovic, B., Arifi, F., & Bjelica, D. (2017). Standing Height and its Estimation Utilizing Foot Length Measurements in Adolescents from Western Region in Kosovo. *Sport Mont*, 15(3), 3-7. doi: 10.26773/smj.2017.10.001
- Popovic, S., Bjelica, D., Georgiev, G., Krivokapic, D., & Milasinovic, R. (2016). Body Height and its Estimation Utilizing Arm Span Measurements in Macedonian Adults. *Anthropologist*, 24(3), 737-745.
- Quanjer, P. H., Capderou, A., Mazocioglu, M. M., Aggarwal, A., Popovic, S., Datta Banik, S., Tayie, F. A. K., Golshan, M., Ip, M. S. M., & Zelter, M. (2014). All-age relationship between arm span and height in different ethnic groups. *European Respiratory Journal*, 44(4), 905-912.
- Reeves, S. L., Varakamin, C., & Henry, C. J. (1996). The relationship between arm-span measurements and height with special reference to gender and ethnicity. *European Journal of Clinical Nutrition*, 50(6), 398-400.
- Starc, G., Popović, S., Đorđić, V., Ostojčić, S., Musić Milanović, S., Kujundžić, E., Spiroski, I., Đurić, S., Mašanović, B., Sember, V., & Leskošek, B. (2019). Differences in body height between the contemporary Western Balkan children and the WHO growth references core sample. *Anthropological Notebook*, 25(3), 55-67.
- Steele, M. F., & Chenier, T. C. (1990). Arm-span, height and age in black and white women. *Annals of Human Biology*, 17(6), 533-541.
- Szollosi, E. (1998). Secular trend in Debrecen university students (in Hungarian). *Anthropologiai Közlemények*, 39, 43-51.
- Ter Goon, D., Toriola, A. T., Musa, D. I., & Akusu, S. (2011). The relationship between arm span and stature in Nigerian adults. *Kinesiology*, 43(1), 38-43.
- Wronka, I., & Pawlinska-Chmara, R. (2009). Childhood environment and adult height among Polish university students. *Collegium Antropologicum*, 33(4), 1039-1045.